

JBP

Journal of Biochemicals and Phytomedicine

eISSN: 2958-8561



Fieldwork and literature review to identify fruits with antidiabetic properties in the Jaffna District, Sri Lanka

Saravanan Vivekanandarajah ^{1,2*} , Pholtan Rajamanoharan ^{3,4} , Vinujan Shanmugalingam ⁵

¹ KnowledgeLink Group, Inc., Waltham, MA 02451, USA

² Poigai Institute, Batticaloa 30000, Sri Lanka

³ Provincial Herbal Garden Management Center, Trincomalee 31000, Sri Lanka

⁴ District Siddha Ayurvedic Hospital, Nilaveli, Trincomalee 31010, Sri Lanka

⁵ Palmyra Research Institute, Jaffna 40000, Sri Lanka

ARTICLE INFO

Article Type:

Research

Article History:

Received: 23 Oct 2024

Revised: 29 Nov 2024

Accepted: 08 Dec 2024

Available online: 31 Dec 2024

Keywords:

Diabetes,
Sri Lanka,
Jaffna,
Myrtaceae,
Garcinia mangostana

*Corresponding author:

E-mail: vivekanandarajahs@yahoo.co.uk

ABSTRACT

Introduction: Antidiabetic medications are often associated with side effects such as mood changes, dizziness, and breathing difficulties. Studies have suggested that fruit consumption may contribute to the prevention and management of diabetes. This study aimed to document the fruits sold in the Jaffna District, Sri Lanka, through fieldwork conducted at markets, fruit kiosks, shops, superstores, and street vendors and, to determine the antidiabetic potential of the identified fruits.

Methods: To conduct this study, researchers visited key markets and local kiosks at least three times to document the presence of fruits, identify different species, and observe how they are used in local diets. The antidiabetic potential of the identified fruits was assessed by reviewing scientific evidence from published studies in electronic databases, including Web of Science, PubMed, Scopus, and ScienceDirect, up to September 2023.

Results: A total of 103 fruit species belonging to 43 families were identified, 9% of which are utilized in antidiabetic preparations in Sri Lankan Siddha Medicine. Among the identified fruits, 52% demonstrated in vivo evidence, 22% were supported by in vitro studies, and 20% had clinical evidence. Additionally, 22 active compounds were isolated from these fruits.

Conclusion: This study serves as a valuable resource for future research on the antidiabetic potential of fruits in the Jaffna District. Further investigations are needed to fully explore their therapeutic applications in diabetes management.

Please cite this paper as:

Vivekanandarajah S, Rajamanoharan PH, Shanmugalingam V. Fieldwork and literature review to identify fruits with antidiabetic properties in the Jaffna District, Sri Lanka. Journal of Biochemicals and Phytomedicine. 2024; 3(2): 83-94. doi: 10.34172/jbp.2024.22.

Introduction

In 2021, roughly 537 million people between the ages of 20 and 79 were living with diabetes. This number is expected to rise to 700 million by 2045. The majority (90%) of the people have type 2 diabetes, while 10% have type 1 diabetes (International Diabetes Federation, 2023). Furthermore, approximately 79% have diabetes reside in low-income and middle-income nations. One in five adults (above 65 years) have diabetes in the world. Unfortunately, around 232 million people (half of those with diabetes) are undiagnosed. This can lead to serious complications, such as cardiovascular diseases, nerve damage, and kidney failure (World Health Organization, 2023). So far, 4.2 million people have died from diabetes, and diabetes causes 1.6 million deaths each year. In addition, at the minimum of 760 billion US dollars spent for diabetes in 2019. It was nearly 10% of the total adult healthcare expenditure (International Diabetes Federation, 2023). Approximately 1.1 million children are diagnosed with type 1 diabetes in the world. In addition, one in six live births (20 million) is affected by gestational diabetes. Interestingly, 374 million people are at risk of getting type 2 diabetes (World Health Organization, 2023).

Increased blood glucose concentration in the body is called diabetes, and it is a metabolic and chronic illness. Diabetes leads to complications of kidney failure, stroke, blindness, heart attack, amputation, etc. (World Health Organization, 2023). Also, the three types of diabetes are type 1 diabetes, type 2 diabetes, and gestational diabetes. Risk factors for diabetes include obesity, physical inactivity, poor diet, and a family history of the disease (European Association for the Study of Diabetes, 2023). Some behaviors like obesity, overweight, unhealthy food, and physical inactivity can increase the risk factors. Anyhow, taking key actions such as healthy eating, avoiding unnecessary weight gain, being physically dynamic, monitoring glucose concentrations, and following medical advice could reduce the risks of getting diabetes (Mayo Foundation for Medical Education and Research, 2023). Type 1 diabetes is managed by injecting insulin into the body. Type 2 diabetes is managed by taking the medications like Metformin and Sulfonylureas (Glibenclamide, Gliclazide, Glimepiride, Glipizide, and Tolbutamide) (International Diabetes Federation, 2023). These medications cause common side effects including a metallic taste, constipation, dark urine, diarrhea, easy bleeding, fever, headache, sore throat, itching, swelling, loss of appetite, mood changes, nausea, rash, yellowing eyes, seizures, weakness, severe dizziness, stomach pain, stomach upset, hands swelling, feet swelling, trouble breathing, unusual tiredness, weight gain, vomiting, weight gain, and yellowing skin (American

Diabetes Association, 2023; WebMD LLC, 2023).

Given these concerning global statistics, it is critical to explore alternative, low-cost, and accessible treatments. One promising area of research is the potential role of fruits, which have shown promise in managing blood sugar levels. Fruit consumption showed prevention and management of diabetes in researches (Feskens et al., 1995; Van Dam et al., 2002). Several research articles were published regarding the antidiabetic activities of various fruits. For example, Intake of fruit juice and incidence of type 2 diabetes: A systematic review and meta-analysis, Potential health benefits of fruits and vegetables: Epic inspite glycemia Food groups in dietary prevention of type 2 diabetes, Fruit and vegetable intake and incidence of type 2 diabetes mellitus: Systematic review and meta-analysis, Prevention of metabolic disorders: Fruits (including fruit sugars) vs. vegetables (Carter et al., 2010; Kuzma et al., 2017; Xi et al., 2014). Polyphenols found in fruits are antioxidants that have defensive properties (Anderson et al., 2004). Fruits such as berries, apples, and citrus are rich in polyphenols and fiber, which have been shown to improve insulin sensitivity and reduce blood sugar levels. Fruits have insulin-like and enhancing insulin secretion effects (Survay et al., 2010; Wedick et al., 2012). Polyphenols in fruits may help manage diabetes by improving insulin sensitivity, inhibiting carbohydrate-digesting enzymes like α -amylase and α -glucosidase, and reducing oxidative stress. These compounds have shown promise in regulating postprandial blood sugar levels, making them key targets for future diabetes treatments.

As mentioned before, biomedicine medications and treatments cause adverse side effects, and they are expensive. Therefore, it is essential to identify natural and cost-effective treatments that can be used to manage diabetes. Hence, this research is crucial because, while numerous studies have explored the antidiabetic potential of fruits globally, few have focused on the fruits widely available in Sri Lanka, particularly in the Jaffna District. Understanding these local fruits could offer an affordable and accessible solution for diabetes management. It assesses the levels of scientific antidiabetic evidence available for the documented fruits. This work benefits the public to manage or prevent diabetes naturally and economically by consuming fruits with antidiabetic properties. Moreover, it is useful for the researchers to study the possible fruits to identify potential antidiabetic active extracts and compounds. These compounds may serve as potential candidates for the development of future antidiabetic drugs.

Materials and methods

The Study Region

This study was performed in the Jaffna District in the Northern Province of Sri Lanka (Figure 1). Jaffna District has a 1,025 km² area and had a population of 624,179 in 2017. The majority of the population is Sri Lankan Tamil, and the Tamil language is mostly spoken in the study region (Annual Performance and Accounts Report - Jaffna District, 2017).

Data Collection

This study was conducted from January 2019 to June 2023. Field visits were conducted at least three times to key markets and local kiosks to document the availability of fruits, identify species, and observe their use in local diets. This process helped ensure that only widely available fruits were included in the study. The authors spent at least two hours in each visit to each market and the surrounding areas.

The Jaffna District, located in the northern part of Sri Lanka, is characterized by its tropical climate and diverse agricultural practices. This area is home to a variety of fruits that may possess untapped potential for managing diabetes, making it an ideal location for this study. The main markets in the Jaffna district visited and the latitudes and longitudes of each location are given below:

1. Jaffna Town Market (09°39'55.53" N, 80°00'31.12" E)
2. Thirunelveli Market (09°41'19" N, 80°01'40.8" E)
3. Chavakachcheri Market (09°39'32.32" N, 80°09'44.11" E)
4. Kodikamam Market (09°40'57.04" N, 80°13'15.79" E)
5. Point Pedro Market (09°49'29.75" N, 80°14'10.68" E)
6. Nellyadi Market (09°48'00.75" N, 80°11'59.96" E)
7. Pandatharippu Market (09°46'23.89" N, 79°58'16.45" E)
8. Maruthanarmadam Market (09°43'46.37" N, 80°01'22.41" E)
9. Velanai Market (09°37'52.25" N, 79°53'46.73" E)
10. Sankanai Market (09°44'55.52" N, 79°58'13.03" E)




Figure 1: The study region and the fieldworks carried out main markets based on Survey Department of Sri Lanka (2014).

Fruit Identification

All the fruits available for sale at each location were identified and confirmed by Dr. Pholtan Rajamanoharan (Provincial Herbal Garden Management Center, Trincomalee 31000, Sri Lanka).

Voucher Specimens

Voucher specimens of the identified fruit species locally available were collected in the Jaffna District from January 2019 to June 2023. Dr. Fultaan

Rajamanoharan, a botanist with over 20 years of experience in plant taxonomy, confirmed the botanical identification of each species. This ensured that the species included in the study were accurately classified. Plant specimens were carefully preserved in the herbarium, with each sample stored in silica gel to prevent degradation. These samples are available for further study and verification at the Provincial Herbal Garden in Trincomalee. All the

scientific names and families of the identified fruit plant species were validated using Kew Science (2024) and Global Biodiversity Information Facility (2024).

Data Analysis

Data were analyzed using a combination of qualitative and quantitative methods, comparing the identified species with existing databases such as the Kew Science database and published literature. Statistical analyses were conducted to identify significant patterns in the occurrence and medicinal properties of the plants. A record that includes the scientific name, family, Tamil name, and herbarium voucher specimen number (if applicable) of all the identified and confirmed fruit species in the fieldwork was created. Then they were compared with the fruits used to treat diabetes in Sri Lankan Siddha Medicine using previously published articles. For example, (Vivekanandarajah and Jeyaseelan 2020; Vivekanandarajah and Rajamanoharan 2021; Vivekanandarajah et al., 2015, 2016, 2017, 2018; Vivekanandarajah, 2021, 2021a, 2021b (Part 1, 2, 3), 2021c), and Rajamanoharan (2014; 2016). Other ethnobotanical uses analysis was performed using the published works in Sri Lanka. For example, Jayaweera (1980; 1981; 1982). In addition, the electronic databases namely, Web of Science, PubMed, Scopus, SpringerLink, ScienceDirect, Wiley Online Library, Taylor & Francis Online, Mary Ann Liebert Inc. Publishers, Wolters Kluwer Medknow Publications, Thieme Medical Publishers, Hindawi Limited, Directory of Open Access Journals (DOAJ), Multidisciplinary Digital Publishing Institute (MDPI), Walter de Gruyter GmbH, BioMed Central Ltd., SAGE Publishing, JSTOR, Public Library of Science (PLOS), and Frontiers Media S.A. were used to identify the published relevant articles until September 2023. Plant species mentioned in American Herbal Pharmacopoeia-Verified Botanical Reference Materials (2023), American Herbal Pharmacopoeia: Botanical Pharmacognosy-Microscopic Characterization of Botanical Medicines (2016), European Medicines Agency's Committee on Herbal Medicinal Products (2023), World Health Organization (WHO) Monographs on Selected Medicinal Plants - Volumes 1 to 4 (1999; 2004; 2007; 2009), and African Herbal Pharmacopoeia (2010) was very well studied for global plant species. Hence, these plant species were excluded from identifying the antidiabetic activities-related articles. The scientific name of each plant species was used as a search term. Only identified fruit species showed antidiabetic activities were considered in this work. Then, the available levels of scientific evidence of antidiabetic activity of each species were assessed.

Results

Most of the fruits were identified in Jaffna Town and Thirunelveli markets, followed by, Kodikamam, Chavakachcheri, Point Pedro, and Nelliyadi markets. On one hand, *Aegle marmelos*, *Anacardium occidentale*, *Ananas comosus*, *Annona cherimola*, *A. muricata*, *A. reticulata*, *A. squamosa*, *Artocarpus heterophyllus*, *Averrhoa bilimbi*, *A. carambola*, *Borassus flabellifer*, *Carica papaya*, *Carissa carandas*, *Citrullus lanatus*, *Citrus × aurantium*, *C. maxima*, *C. medica*, *Cocos nucifera*, *Cordia dichotoma*, *Cucumis melo*, *Donella lanceolata*, *Ficus racemosa*, *Fragaria × ananassa*, *Garcinia mangostana*, *G. quaesita*, *G. zeylanica*, *Lansium domesticum*, *Limonia acidissima*, *Malus domestica*, *Mangifera indica*, *Manilkara hexandra*, *Musa × paradisiaca*, *Nephelium lappaceum*, *Passiflora edulis*, *Phoenix dactylifera*, *P. pusilla*, *Phyllanthus acidus*, *P. emblica*, *Psidium cattleianum*, *P. guajava*, *Punica granatum*, *Schleichera oleosa*, *Selenicereus undatus*, *Syzygium cumini*, *S. malaccense*, and *Vitis vinifera* fruits were widely identified and available in all visited markets. On the other hand, fruit species namely, *Actinidia chinensis*, *Antidesma bunius*, *Atalantia ceylanica*, *Baccaurea motleyana*, *Bridelia retusa*, *Calophyllum calaba*, *Canthium coromandelicum*, *Careya arborea*, *Carissa spinarum*, *Citrus japonica*, *Dillenia retusa*, *Dimocarpus longan*, *Diospyros malabarica*, *Elaeocarpus serratus*, *Erythroxylum moonii*, *Flacourtia indica*, *F. jangomas*, *Gmelina arborea*, *Grewia tiliifolia*, *Huberanthia korinti*, *Lantana camara*, *Mangifera zeylanica*, *Memecylon intermedium*, *Mimusops elengi*, *Morus alba*, *Opuntia stricta*, *Passiflora foetida*, *Phyllanthus reticulatus*, *Pouteria campechiana*, *Pyrus communis*, *P. pyriformis*, *Salacia chinensis*, *Sandoricum koetjape*, *Sonneratia caseolaris*, *Synsepalum dulcificum*, *Syzygium aqueum*, *S. caryophyllatum*, *S. nervosum*, *Syzygium samarangense*, *Tarenna asiatica*, *Vateria copallifera*, *Ziziphus linnaei*, and *Z. oenopolia* were rarely identified and available.

A total of 103 fruit species from 43 families were identified and documented in this work (Table 1). Most of the fruits belong to *Syzygium*, followed by *Citrus*, *Annona*, and *Ziziphus* species. Most of the fruits were from the *Myrtaceae* family, followed by *Rutaceae*, *Sapotaceae*, *Phyllanthaceae*, and *Annonaceae*.

Discussion

Many of the identified fruits were from food plants. Nine fruit species including *A. marmelos*, *A. occidentale*, *A. carambola*, *B. flabellifer*, *L. acidissima*, *M. paradisiaca*, *P. dactylifera*, *P. emblica*, and *P. granatum* are used in Sri Lankan Siddha Medicine antidiabetic preparations (Vivekanandarajah and Jeyaseelan 2020; Vivekanandarajah and Rajamanoharan 2021; Vivekanandarajah et al., 2015, 2016, 2017, 2018; Vivekanandarajah, 2021, 2021a, 2021b, 2021c).

Levels of Scientific Evidence of Identified Fruit Species

A sum of 26 (25%) fruit species was very well studied and globally distributed. Hence, a literature review of 77 fruit species revealed that 25 fruit species (32%) have antidiabetic scientific evidence. Most numbers of the fruit species with antidiabetic properties were from the *Annonaceae* family, followed by *Rutaceae* and *Sapotaceae*.

Fruit Species Had in vitro Antidiabetic Scientific Evidence

A total of seven fruits (28%) had *in vitro* antidiabetic scientific evidence as to the highest level (Table 2). *A. cherimola* and *S. dulcis* had the majority of evidence (Vasarri et al., 2020; Galarce-Bustos et al., 2019; Hossain et al., 2008; Mohamed Yunus et al., 2021). No antidiabetic compound was isolated from these fruit species.

Table 1: Fruit species sold in markets in jaffna district

Scientific name	Family	Tamil	Herbarium voucher specimen identification
<i>Actinidia chinensis</i> Planch.	<i>Actinidiaceae</i>	Kiwi	NA
<i>Aegle marmelos</i> (L.) Corrêa	<i>Rutaceae</i>	Vilvai	PR-1
<i>Anacardium occidentale</i> L.	<i>Anacardiaceae</i>	Munthirihai	PR-10
<i>Ananas comosus</i> (L.) Merr.	<i>Bromeliaceae</i>	Annaasi	NA
<i>Annona cherimola</i> Mill.	<i>Annonaceae</i>	Parangi Annamunnaa	PR-26
<i>Annona muricata</i> L.	<i>Annonaceae</i>	Seeththaa	PR-37
<i>Annona reticulata</i> L.	<i>Annonaceae</i>	Iraama Seeththaa	NA
<i>Annona squamosa</i> L.	<i>Annonaceae</i>	Annamunnaa	PR-52
<i>Antidesma bunius</i> (L.) Spreng.	<i>Phyllanthaceae</i>	Naalaithali	NA
<i>Artocarpus heterophyllus</i> Lam.	<i>Moraceae</i>	Palaa	PR-55
<i>Atalantia ceylanica</i> (Arn.) Oliv.	<i>Rutaceae</i>	Kurunthu	PR-56
<i>Averrhoa bilimbi</i> L.	<i>Oxalidaceae</i>	Vilimbi	PR-2
<i>Averrhoa carambola</i> L.	<i>Oxalidaceae</i>	Thamaraththai	PR-3
<i>Baccaurea motleyana</i> (Müll.Arg.) Müll.Arg.	<i>Phyllanthaceae</i>	Moottipuli	NA
<i>Borassus flabellifer</i> L.	<i>Arecaceae</i>	Panai	PR-4
<i>Bridelia retusa</i> (L.) A.Juss.	<i>Phyllanthaceae</i>	Mulvengai	NA
<i>Calophyllum calaba</i> L.	<i>Calophyllaceae</i>	Manjatpunnai	PR-5
<i>Canthium coromandelicum</i> (Burm.f.) Alston	<i>Rubiaceae</i>	Kaarai	PR-6
<i>Careya arborea</i> Roxb.	<i>Lecythidaceae</i>	Aayimaa	NA
<i>Carica papaya</i> L.	<i>Caricaceae</i>	Pappaasi	PR-7
<i>Carissa carandas</i> L.	<i>Apocynaceae</i>	Perungkalaa	PR-8
<i>Carissa spinarum</i> L.	<i>Apocynaceae</i>	Kalaa	PR-9
<i>Chrysophyllum cainito</i> L.	<i>Sapotaceae</i>	Seemai Laavul	NA
<i>Citrullus lanatus</i> (Thunb.) Matsum. & Nakai	<i>Cucurbitaceae</i>	Thatpoosani	NA
<i>Citrus × aurantium</i> L.	<i>Rutaceae</i>	Then Thodai	PR-11
<i>Citrus × limon</i> (L.) Osbeck	<i>Rutaceae</i>	Malaialumichchai	NA
<i>Citrus japonica</i> Thunb.	<i>Rutaceae</i>	Kaattuth Thodai	PR-12
<i>Citrus maxima</i> (Burm.) Merr.	<i>Rutaceae</i>	Naaraththai	NA
<i>Citrus medica</i> L.	<i>Rutaceae</i>	Elumichchai	PR-13
<i>Cocos nucifera</i> L.	<i>Arecaceae</i>	Thennai	PR-14
<i>Cordia dichotoma</i> G.Forst.	<i>Boraginaceae</i>	Naruvili	PR-15
<i>Cucumis melo</i> L.	<i>Cucurbitaceae</i>	Waththhai	PR-16
<i>Cynometra cauliflora</i> L.	<i>Fabaceae</i>	Naminam	NA

<i>Dialium ovoideum</i> Thwaites	<i>Fabaceae</i>	Kaattuppuli	NA
<i>Dillenia retusa</i> Thunb.	<i>Dilleniaceae</i>	Naaiththekku	NA
<i>Dimocarpus longan</i> Lour.	<i>Sapindaceae</i>	Nurai	NA
<i>Diospyros malabarica</i> (Desr.) Kostel.	<i>Ebenaceae</i>	Panichchai	PR-17
<i>Donella lanceolata</i> (Blume) Aubrév.	<i>Sapotaceae</i>	Ilaavul	PR-18
<i>Drypetes sepiaria</i> (Wight & Arn.) Pax & K.Hoffm.	<i>Putranjivaceae</i>	Weerai	PR-19
<i>Durio zibethinus</i> L.	<i>Malvaceae</i>	Mulnaari	NA
<i>Elaeocarpus serratus</i> L.	<i>Elaeocarpaceae</i>	Weralu	NA
<i>Erythroxylum moonii</i> Hochr.	<i>Erythroxylaceae</i>	Sirusemmanaththai	NA
<i>Ficus racemosa</i> L.	<i>Moraceae</i>	Aththi	PR-20
<i>Flacourtia indica</i> (Burm.f.) Merr.	<i>Salicaceae</i>	Kaattukkalai	PR-21
<i>Flacourtia inermis</i> Roxb.	<i>Salicaceae</i>	Lovi	NA
<i>Flacourtia jangomas</i> (Lour.) Raeusch.	<i>Salicaceae</i>	Vaiyangkaarai	NA
<i>Fragaria</i> × <i>ananassa</i> (Duchesne ex Weston) Duchesne ex Rozier	<i>Rosaceae</i>	Strawberry	NA
<i>Garcinia mangostana</i> L.	<i>Clusiaceae</i>	Mangusththaan	NA
<i>Garcinia quaesita</i> Pierre	<i>Clusiaceae</i>	Korakkaappuli	NA
<i>Garcinia zeylanica</i> Roxb.	<i>Clusiaceae</i>	Kattuk Korakkaappuli	NA
<i>Gmelina arborea</i> Roxb. ex Sm.	<i>Lamiaceae</i>	Kumil	PR-22
<i>Grewia tiliifolia</i> Vahl	<i>Malvaceae</i>	Thavittai	PR-23
<i>Huberantha korinti</i> (Dunal) Chaowasku	<i>Annonaceae</i>	Karuvalli	NA
<i>Lansium domesticum</i> Corrêa	<i>Meliaceae</i>	Kadukuda	NA
<i>Lantana camara</i> L.	<i>Verbenaceae</i>	Arisimalar	PR-24
<i>Limonia acidissima</i> L.	<i>Rutaceae</i>	Vilaa	PR-25
<i>Malus domestica</i> (Suckow) Borkh.	<i>Rosaceae</i>	Apple	NA
<i>Mangifera indica</i> L.	<i>Anacardiaceae</i>	Maa	PR-27
<i>Mangifera zeylanica</i> (Blume) Hook.f.	<i>Anacardiaceae</i>	Kaattu Maa	NA
<i>Manilkara hexandra</i> (Roxb.) Dubard	<i>Sapotaceae</i>	Paalai	PR-28
<i>Manilkara zapota</i> (L.) P.Royen	<i>Sapotaceae</i>	Seemayilupai	PR-29
<i>Memecylon intermedium</i> Blume	<i>Melastomataceae</i>	Kaayaa	PR-30
<i>Mimusops elengi</i> L.	<i>Sapotaceae</i>	Mahilam	PR-31
<i>Morus alba</i> L.	<i>Moraceae</i>	Mayirkkottichchedi	PR-32
<i>Musa</i> × <i>paradisiaca</i> L.	<i>Musaceae</i>	Vaalai	PR-33
<i>Nephelium lappaceum</i> L.	<i>Sapindaceae</i>	Irambuttaan	NA
<i>Opuntia stricta</i> (Haw.) Haw.	<i>Cactaceae</i>	Naahathaali	PR-34
<i>Passiflora edulis</i> Sims	<i>Passifloraceae</i>	Kodiththodai	PR-35
<i>Passiflora foetida</i> L.	<i>Passifloraceae</i>	Sirupoonaiikkaali	PR-36
<i>Persea americana</i> Mill.	<i>Lauraceae</i>	Vennai Maram	NA
<i>Phoenix dactylifera</i> L.	<i>Arecaceae</i>	Pereechchai	NA
<i>Phoenix pusilla</i> Gaertn.	<i>Arecaceae</i>	Eechchai	PR-38
<i>Phyllanthus acidus</i> (L.) Skeels	<i>Phyllanthaceae</i>	Arunelli	PR-39
<i>Phyllanthus emblica</i> L.	<i>Phyllanthaceae</i>	Nelli	PR-40
<i>Phyllanthus reticulatus</i> Poir.	<i>Phyllanthaceae</i>	Sempoolaa	PR-41

<i>Pouteria campechiana</i> (Kunth) Baehni	<i>Sapotaceae</i>	Kaattilaavul	PR-42
<i>Psidium cattleyanum</i> Sabine	<i>Myrtaceae</i>	Kilokkoyyaa	NA
<i>Psidium guajava</i> L.	<i>Myrtaceae</i>	Koyyaa	PR-43
<i>Psidium guineense</i> Sw.	<i>Myrtaceae</i>	Pulikkoyyaa	NA
<i>Punica granatum</i> L.	<i>Lythraceae</i>	Maathulai	PR-44
<i>Pyrus communis</i> L.	<i>Rosaceae</i>	Salvaagu	NA
<i>Pyrus pyrifolia</i> (Burm.f.) Nakai	<i>Rosaceae</i>	Seenach Chalvaahu	NA
<i>Salacia chinensis</i> L.	<i>Celastraceae</i>	Mallivembu	NA
<i>Sandoricum koetjape</i> (Burm.f.) Merr.	<i>Meliaceae</i>	Sevvai	NA
<i>Schleichera oleosa</i> (Lour.) Oken	<i>Sapindaceae</i>	Kumbaththiri	NA
<i>Selenicereus undatus</i> (Haw.) D.R.Hunt	<i>Cactaceae</i>	Dragon	NA
<i>Sonneratia caseolaris</i> (L.) Engl.	<i>Lythraceae</i>	Kinnai	PR-45
<i>Spondias dulcis</i> Parkinson	<i>Anacardiaceae</i>	Ambirala	PR-46
<i>Synsepalum dulcificum</i> (Schumach. & Thonn.) Daniell	<i>Sapotaceae</i>	Atputham	NA
<i>Syzygium aqueum</i> (Burm.f.) Alston	<i>Myrtaceae</i>	Neerchchambanaaval	NA
<i>Syzygium caryophyllatum</i> (L.) Alston	<i>Myrtaceae</i>	Sirunaaval	PR-47
<i>Syzygium cumini</i> (L.) Skeels	<i>Myrtaceae</i>	Naaval	PR-48
<i>Syzygium jambos</i> (L.) Alston	<i>Myrtaceae</i>	Sambanaaval	PR-49
<i>Syzygium malaccense</i> (L.) Merr. & L.M.Perry	<i>Myrtaceae</i>	Periya Sambanaaval	NA
<i>Syzygium nervosum</i> A.Cunn. ex DC.	<i>Myrtaceae</i>	Naahai	NA
<i>Syzygium samarangense</i> (Blume) Merr. & L.M.Perry	<i>Myrtaceae</i>	Neerkkumali	NA
<i>Tarenna asiatica</i> (L.) Kuntze ex K.Schum.	<i>Rubiaceae</i>	Tharani	NA
<i>Vateria copallifera</i> (Retz.) Alston	<i>Dipterocarpaceae</i>	Kungiliyampinai	NA
<i>Vitis vinifera</i> L.	<i>Vitaceae</i>	Thiraatchai	NA
<i>Ziziphus jujuba</i> Mill.	<i>Rhamnaceae</i>	Ilanthai	PR-50
<i>Ziziphus linnaei</i> M.A.Lawson	<i>Rhamnaceae</i>	Soorai	PR-51
<i>Ziziphus mauritiana</i> Lam.	<i>Rhamnaceae</i>	Perilanthai	PR-53
<i>Ziziphus oenopolia</i> (L.) Mill.	<i>Rhamnaceae</i>	Soorayilanthai	PR-54

Table 2: Fruits have *in vitro* scientific evidence

Scientific names	Extract compound	Assay / model / human subject	Citation
<i>Annona cherimola</i>	Ethanol	Non-enzymatic glycation of human serum albumin inhibitory; α -Glucosidase inhibitory	(Vasarri et al., 2020)
<i>Annona cherimola</i>	NA	α -Glucosidase inhibitory	(Galarce-Bustos et al., 2019)
<i>Annona muricata</i>	Aqueous	α -Amylase inhibitory; α -Glucosidase inhibitory	(Adefegha et al., 2015)
<i>Chrysophyllum cainito</i>	Ethanol	α -Glucosidase inhibitory	(Ningsih et al., 2020)
<i>Phoenix pusilla</i>	Ethanol	α -Amylase inhibitory; α -Glucosidase inhibitory	(Sankar and Shoba, 2015)
<i>Phyllanthus acidus</i>	NA	α -Glucosidase inhibitory	(Suliaman and Ooi, 2014)
<i>Phyllanthus acidus</i>	Ethanol	α -Glucosidase inhibitory	(Zulaikha et al., 2018)
<i>Spondias dulcis</i>	Ethanol	α -Amylase inhibitory; α -Glucosidase inhibitory	(Hossain et al., 2008)
<i>Spondias dulcis</i>	Ethanol	α -Glucosidase inhibitory	(Yunus et al., 2021)
<i>Syzygium caryophyllatum</i>	Ethanol	α -Amylase inhibitory	(Wathsara et al., 2020)

Table 3: Fruits have in vivo scientific evidence

Scientific names	Extract/ compounds	Assay / model / human subject	Reference
<i>Annona squamosa</i>	NA	Alloxan-induced diabetic	(Gupta et al., 2005)
<i>Averrhoa bilimbi</i>	Aqueous	Streptozotocin-induced diabetic	(Kurup and Mini, 2016), (Kurup and Mini, 2017)
<i>Carissa carandas</i>	Methanol	Alloxan-induced diabetic	(Itankar et al., 2011)
<i>Citrus maxima</i>	Ethanol	Alloxan-induced diabetic	(Sci et al., 2020)
<i>Citrus medica</i>	NA	NS	(Peng et al., 2009)
<i>Ficus racemosa</i>	Ethanol	Type 1 diabetic; Type 2 diabetic	(Trinh et al., 2017)
<i>Ficus racemosa</i>	α -Amyrin acetate	Streptozotocin-induced diabetic	(Narender et al., 2009)
<i>Gmelina arborea</i>	Butanol, Ethanol, Ethyl acetate, Petroleum ether	Alloxan-induced diabetic	(Nayak et al., 2012)
<i>Limonia acidissima</i>	NA	Fluoride-induced diabetic	(Vasant and Narasimhacharya, 2013)
<i>Manilkara zapota</i>	Aqueous	Normal	(Barbalho et al., 2015)
<i>Synsepalum dulcificum</i>	Ethanol	Alloxan-induced diabetic	(Haddad et al., 2020)
<i>Synsepalum dulcificum</i>	NA	Fructose-rich chow-fed-induced insulin resistant	(Jang et al., 2008)
<i>Syzygium samarangense</i>	Vescalagin	High-fructose diet-induced diabetic	(Shen and Chang, 2013), (Huang et al., 2016)

Table 4: Fruits have clinical scientific evidence

Scientific names	Extract / compound	Human subject	Reference
<i>Borassus flabellifer</i>	NA	Type 2 diabetic	(Rahman et al., 2020)
<i>Garcinia mangostana</i>	α -Mangostin, γ -Mangostin	Obese female	(Watanabe et al., 2018)
<i>Mangifera indica</i>	NA	Obese	(Evans et al., 2014)
<i>Passiflora edulis</i>	NS	Type 2 diabetic	(De Queiroz et al., 2012)

Conclusion

Fruits are one of the most important parts of daily food. The primary goal of this study is to evaluate the scientific evidence for the antidiabetic properties of fruits commonly found in the Jaffna District. As mentioned above, many of the fruits have no antidiabetic scientific evidence. Hence, more studies should be conducted to identify the antidiabetic potentials of these fruits. Furthermore, priority should be provided to the widely available fruits in these future studies. Thus, these fruits will be useful to manage diabetes with fewer side effects in an economical way. Widely available fruits also have easier access, and they are affordable. The identified fruits in the study contain several antidiabetic active compounds. Therefore, further researches should be carried out to study these compounds in more advanced models to create more scientific evidence. This study identified, documented, and assessed the antidiabetic activities of fruits currently sold in the

Jaffna District in Sri Lanka. This research provides a foundation for future studies on locally available fruits and their potential role in diabetes management.

Declarations

Conflict of interest

The authors declare that there is no conflict of interest.

Acknowledgments

The authors are grateful to their family members for their support in delivering this work.

Consent for publications

All authors have read and approved the manuscript for publication.

Funding/support

This work received no funding.

Authors' contributions

SV conceptualized and designed the the study. PR and VS coppedelaborated in theon data curation and carried out the study. All authors had the same role in carrng outstudy implementation. All authors contributed equally to conducting the literature search and wridrafting the first draft of thetheinitial manuscript. All authors reaviewed and approved the final manuscript for publication.

Ethical Considerations

Ethical issues (including plagiarism, data fabrication, double publication and submission, redundancy) have been completely looked into by the author.

References

- Abdallah HM, El-Bassossy H, Mohamed GA, El-Halawany AM, Alshali KZ, Banjar ZM. Phenolics from *Garcinia mangostana* inhibit advanced glycation end products formation: Effect on Amadori products, cross-linked structures and protein thiols. *Molecules*. 2016;21(2):251. doi:10.3390/molecules21020251.
- Abdallah HM, El-Bassossy H, Mohamed GA, El-Halawany AM, Alshali KZ, Banjar ZM. Mangostanaxanthones III and IV: Advanced glycation end-product inhibitors from the pericarp of *Garcinia mangostana*. *Journal of Natural Medicines*. 2017;71(2):216–226. doi:10.1007/s11418-016-1051-8.
- Adefegha SA, Oyeleye SI, Oboh G. Distribution of phenolic contents, antidiabetic potentials, antihypertensive properties, and antioxidative effects of soursop (*Annona muricata* L.) fruit parts in vitro. *Biochemistry Research International*. 2015;2015:347673. doi:10.1155/2015/347673.
- American Diabetes Association. Diabetes. Available at: <https://www.diabetes.org/diabetes>. Accessed October 5, 2023.
- American Herbal Pharmacopoeia. Botanical Reference Standards. Available at: http://www.herbal-ahp.org/order_online.htm. [Accessed May 17, 2023.]
- American Herbal Pharmacopoeia-Verified Botanical Reference Materials. Available at: <https://herbal-ahp.org/botanical-reference-materials/>. [Accessed May 19, 2023.]
- Anderson RA, Broadhurst CL, Polansky MM, Schmidt WF, Khan A, Flanagan VP, Schoene NW, Graves DJ. Isolation and characterization of polyphenol type-A polymers from cinnamon with insulin-like biological activity. *Journal of Agricultural and Food Chemistry*. 2004;52(6):1475–1481. doi:10.1021/jf035274.
- Annual Performance and Accounts Report - Jaffna District, 2017. Available at: <https://www.parliament.lk/uploads/documents/paperspresented/performance-report-district-secretariat-jaffna-2017.pdf>. [Accessed October 5, 2023.]
- Barbalho SM, Bueno PCDS, Delazari DS, Guiguer EL, Coqueiro DP, Araújo AC, De Souza MDSS, Farinazzi-Machado FMV, Mendes CG, Groppo M. Antidiabetic and antilipidemic effects of *Manilkara zapota*. *Journal of Medicinal Food*. 2015;18(3):385–391. doi:10.1089/jmf.2013.0170.
- Brendler T. *African Herbal Pharmacopoeia*. Port Louis: Association for African Medicinal Plants Standards; 2010.
- Carter P, Gray LJ, Troughton J, Khunti K, Davies MJ. Fruit and vegetable intake and incidence of type 2 diabetes mellitus: Systematic review and meta-analysis. *BMJ*. 2010;341:c4229. doi:10.1136/bmj.c4229.
- De Queiroz A, De Faveri R, Broering MF, Bousfield IT, Goss MJ, Muller SP, Pereira RO, Silva AMO, Machado ID, Quintão NLM, Santin JR. Effects of passion fruit peel flour (*Passiflora edulis* f. *flavicarpa* O. Deg.) in cafeteria diet-induced metabolic disorders. *Journal of Ethnopharmacology*. 2020;250:112482. doi:10.1016/j.jep.2019.112482.
- European Association for the Study of Diabetes. Empower e-learning. Available at: <https://www.easd.org/education/empower-e-learning.html>. [Accessed August 6, 2023.]
- European Medicines Agency. Committee on Herbal Medicinal Products (HMPC). Available at: http://www.ema.europa.eu/ema/index.jsp?curl=pages/about_us/general/general_content_000264.jsp. [Accessed May 4, 2023.]
- European Medicines Agency's Committee on Herbal Medicinal Products. Available at: <https://www.ema.europa.eu/en/committees/committee-herbal-medicinal-products-hmpc>. [Accessed July 9, 2023.]
- Evans SF, Meister M, Mahmood M, Eldoumi H, Peterson S, Perkins-Veazie P, Clarke SL, Payton M, Smith BJ, Lucas EA. Mango supplementation improves blood glucose in obese individuals. *Nutrition and Metabolic Insights*. 2014;7:77–84. doi:10.4137/NMI.S17028.
- Feskens EJ, Virtanen SM, Räsänen L, Tuomilehto J, Stengård J, Pekkanen J, Nissinen A, Kromhout D. Dietary factors determining diabetes and impaired glucose tolerance: A 20-year follow-up of the Finnish and Dutch cohorts of the Seven Countries Study. *Diabetes Care*. 1995;18(8):1104–1112.
- Galarce-Bustos O, Pavón-Pérez J, Henríquez-Aedo K, Aranda M. An improved method for a fast screening of α -glucosidase inhibitors in cherimoya fruit (*Annona cherimola* Mill.) applying effect-directed analysis via high-performance thin-layer chromatography-bioassay-mass spectrometry. *Journal of Chromatography A*. 2019;460415. doi:10.1016/j.chroma.2019.460415.

- Global Biodiversity Information Facility. Available at: <https://www.gbif.org/>. [Accessed June 12, 2023.]
- Gupta RK, Kesari AN, Watal G, Murthy PS, Chandra R, Tandon V. Nutritional and hypoglycemic effect of fruit pulp of *Annona squamosa* in normal healthy and alloxan-induced diabetic rabbits. *Annals of Nutrition and Metabolism*. 2005;49(6):407–413.
- Haddad SG, Mohammad M, Raafat K, Saleh FA. Antihyperglycemic and hepatoprotective properties of miracle fruit (*Synsepalum dulcificum*) compared to aspartame in alloxan-induced diabetic mice. *Journal of Integrative Medicine*. 2020;18(6):514–521. doi:10.1016/j.joim.2020.09.001.
- Hossain SJ, Tsujiyama I, Takasugi M, Islam MA, Biswas RS, Aoshima H. Total phenolic content, antioxidative, anti-amylase, anti-glucosidase, and antihistamine release activities of Bangladeshi fruits. *Food Science and Technology Research*. 2008;14(3):261–268. doi:10.3136/fstr.14.261.
- Huang DW, Chang WC, Wu JSB, Shih RW, Shen SC. Vescalagin from pink wax apple (*Syzygium samarangense* (Blume) Merrill and Perry) alleviates hepatic insulin resistance and ameliorates glycemic metabolism abnormality in rats fed a high-fructose diet. *Journal of Agricultural and Food Chemistry*. 2016;64(5):1122–1129. doi:10.1021/acs.jafc.5b05558.
- Ibrahim SRM, Mohamed GA, Khayat MTA, Ahmed S, Abo-Haded H. α -Amylase inhibition of xanthenes from *Garcinia mangostana* pericarps and their possible use for the treatment of diabetes with molecular docking studies. *Journal of Food Biochemistry*. 2019;e12844. doi:10.1111/jfbc.12844.
- International Diabetes Federation. What is diabetes? Facts & figures. Available at: <https://idf.org/aboutdiabetes/what-is-diabetes/facts-figures.html>. [Accessed June 9, 2023.]
- Itankar PR, Lokhande SJ, Verma PR, Arora SK, Sahu RA, Patil AT. Antidiabetic potential of unripe *Carissa carandas* Linn. fruit extract. *Journal of Ethnopharmacology*. 2011;135(2):430–433. doi:10.1016/j.jep.2011.03.036.
- James R. American Herbal Pharmacopoeia. Botanical pharmacognosy – microscopic characterization of botanical medicines. *Australian Journal of Forensic Sciences*. 2015;48(3):359–361. doi:10.1080/00450618.2015.1121799.
- Jang MH, Piao XL, Kim JM, Kwon SW, Park JH. Inhibition of cholinesterase and amyloid- β aggregation by resveratrol oligomers from *Vitis amurensis*. *Phytotherapy Research*. 2008;22(4):544–549.
- Jayaweera DMA. Medicinal Plants (Indigenous and Exotic) Used in Ceylon (Part 2). A publication of the National Science Council of Sri Lanka. 1980:162–163.
- Jayaweera DMA. Medicinal Plants Used in Ceylon, Vols. 1–4. National Science Council of Sri Lanka, Colombo; 1981.
- Jayaweera DMA. Medicinal Plants Used in Ceylon, first ed. Gunasena MD & Co, Sri Lanka; 1982.
- Karim N, Rahman MA, Changlek S, Tangpong J. Short-time administration of xanthone from *Garcinia mangostana* fruit pericarp attenuates the hepatotoxicity and renotoxicity of type II diabetes mice. *Journal of the American College of Nutrition*. 2020;39:501–510. doi:10.1080/07315724.2019.1696251.
- Kew Science. Plants of the World Online. Available at: <http://plantsoftheworldonline.org/taxon/urn:lsid:ipni.org:names:60474520-2>. [Accessed July 14, 2023.]
- Kurup SB, Mini S. Averrhoa bilimbi fruits attenuate hyperglycemia-mediated oxidative stress in streptozotocin-induced diabetic rats. *Journal of Food and Drug Analysis*. 2016;25(2):360–368.
- Kurup SB, Mini S. Protective potential of Averrhoa bilimbi fruits in ameliorating the hepatic key enzymes in streptozotocin-induced diabetic rats. *Biomedicine & Pharmacotherapy*. 2017;85:725–732.
- Kuzma JN, Schmidt KA, Kratz M. Prevention of metabolic diseases: Fruits (including fruit sugars) vs. vegetables. *Current Opinion in Clinical Nutrition and Metabolic Care*. 2017;20:286–293. doi:10.1097/MCO.0000000000000378.
- Ma Q, Guo Y, Sun L, Zhuang Y. Anti-diabetic effects of phenolic extract from rambutan peels (*Nephelium lappaceum*) in high-fat diet and streptozotocin-induced diabetic mice. *Nutrients*. 2017;9(8):801. doi:10.3390/nu9080801.
- Mayo Foundation for Medical Education and Research. Diabetes Symptoms and Causes. Available at: <https://www.mayoclinic.org/diseases-conditions/diabetes/symptoms-causes/syc-20371444>. [Accessed May 19, 2023.]
- Mohamed Yunus SN, Abas F, Jaafar AH, Azizan A, Zolkeflee NKZ, Ghafar SZA. Antioxidant and α -glucosidase inhibitory activities of eight neglected fruit extracts and UHPLC-MS/MS profile of the active extracts. *Food Science and Biotechnology*. 2021;30:195–208.
- Narender T, Khaliq T, Singh AB, Joshi MD, Mishra P, Chaturvedi JP, Srivastava AK, Maurya R, Agarwal SC. Synthesis of α -amyrin derivatives and their in vivo antihyperglycemic activity. *European Journal of Medicinal Chemistry*. 2009;44(3):1215–1222. doi:10.1016/j.ejmech.2008.09.011.
- Nayak BS, Ellaiah P, Dinda SC. Antibacterial, antioxidant and antidiabetic activities of *Gmelina arborea* Roxb fruit extracts. *International Journal of Green Pharmacy*. 2012;6(3):224–230. doi:10.4103/0973-8258.104937.
- Ningsih IY, Sofyan MD, Prabandari T, Lachtheany V, Hidayat MA. Antioxidant and α -glucosidase inhibitory activities of four types of *Chrysophyllum cainito* L. fruit. *Fabad Journal of Pharmaceutical Sciences*. 2020;45(2):105–115.

- Peng CH, Ker YB, Weng CF, Peng CC, Huang CN, Lin LIY, Peng RY. Insulin secretagogue bioactivity of finger citron fruit (*Citrus medica* L. var. *Sarcodactylis* Hort, Rutaceae). *Journal of Agricultural and Food Chemistry*. 2009;57(19):8812–8819. doi:10.1021/jf902143x.
- Rahman SS, Yasmin N, Kamruzzaman MD, Rezwanaul Islam MD, Rezaul Karim MD, Rouf MA. Anti-hyperglycemic effect of the immature endosperm of sugar palm (*Borassus flabellifer*) fruit on type 2 diabetes mellitus patients – a case study. *Diabetes & Metabolic Syndrome: Clinical Research & Reviews*. 2020;14:1317–1322.
- Rajamanoharan PRS. An ethnobotanical survey of medicinal plants in Sillalai, Jaffna, Northern Province, Sri Lanka. *International Journal of Herbal Medicine*. 2014;1:22–30.
- Rajamanoharan PRS. Existence and survey of medicinal plants of Thiru Palani Murugan Kovil, Asikulam, Vavuniya District, Northern Province, Sri Lanka. *World Journal of Pharmaceutical Research*. 2016;5:161–182. doi:10.20959/wjpr201611-6702.
- Sankar V, Shoba F. Inhibitory effect of Phoenix pusilla unripe fruit on the enzymes, α -amylase and α -glucosidase. *International Journal of PharmTech Research*. 2015;8(1):123–126.
- Sci A, Shaddock APOF, Extract P. Anti-diabetic, anti-hyperlipidemic and hepatoprotective potential of shaddock (*Citrus maxima*) peel extract. *Acta Scientiarum Polonorum. Technologia Alimentaria*. 2020;19(3):271–278.
- Shen SC, Chang WC. Hypotriglyceridemic and hypoglycemic effects of vescalagin from Pink wax apple (*Syzygium samarangense* (Blume) Merrill and Perry cv. Pink) in high-fructose diet-induced diabetic rats. *Food Chemistry*. 2013;136:858–863. doi:10.1016/j.foodchem.2012.08.037.
- Silva DC, Freitas ALP, Pessoa CDS, Paula RCM, Mesquita JX, Leal LKA, Brito GAC, Gonçalves DO, Viana GSB. Pectin from *Passiflora edulis* shows anti-inflammatory action as well as hypoglycemic and hypotriglyceridemic properties in diabetic rats. *Journal of Medicinal Food*. 2011;14:1118–1126. doi:10.1089/jmf.2010.0220.
- Suliaman S, Shaida Fariza Ooi LK. Antioxidant and α -glucosidase inhibitory activities of 40 tropical juices from Malaysia and identification of phenolics from the bioactive fruit juices of *Barringtonia racemosa* and *Phyllanthus acidus*. *Journal of Agricultural and Food Chemistry*. 2014;62(39):9576–9585.
- Survay NS, Ko EY, Upadhyay CP, Jang M, Park SW, Lee DH, Hong SJ. Hypoglycemic effects of fruits and vegetables in hyperglycemic rats for prevention of type-2 diabetes. *Korean Journal of Horticultural Science and Technology*. 2010;28:850–856.
- Trinh BTD, Jäger AK, Staerk D. High-resolution inhibition profiling combined with HPLC-HRMS-SPE-NMR for identification of PTP1B inhibitors from Vietnamese plants. *Molecules*. 2017;22(7):1228. doi:10.3390/molecules22071228.
- Trinh BTD, Staerk D, Jäger AK. Screening for potential α -glucosidase and α -amylase inhibitory constituents from selected Vietnamese plants used to treat type 2 diabetes. *Journal of Ethnopharmacology*. 2016. doi:10.1016/j.jep.2016.03.060.
- Uchida-Maruki H, Inagaki H, Ito R, Kurita I, Sai M, Ito T. Piceatannol lowers the blood glucose level in diabetic mice. *Biological and Pharmaceutical Bulletin*. 2015;38:629–633. doi:10.1248/bpb.b15-00009.
- Van Dam RM, Rimm EB, Willett WC, Stampfer MJ, Hu FB. Dietary patterns and risk for type 2 diabetes mellitus in US men. *Annals of Internal Medicine*. 2002;136:201–209.
- Vasant RA, Narasimhacharya AV. Limonia fruit as a food supplement to regulate fluoride-induced hyperglycemia and hyperlipidemia. *Journal of the Science of Food and Agriculture*. 2013;93(2):422–426. doi:10.1002/jsfa.5762.
- Vasarri M, Barletta E, Vinci S, Ramazzotti M, Francesconi A, Manetti F, Degl'Innocenti A. Annona cherimola Miller fruit as a promising candidate against diabetic complications: An in vitro study and preliminary clinical results. *Foods*. 2020;9:1350. doi:10.3390/foods9101350.
- Vivekanandarajah S. Plants historically used by Sri Lankan Tamils to treat diabetes. Presented at: The International Dr. Safiye Ali Multidisciplinary Studies Congress in Health Sciences; 2021. İzmir, Turkey.
- Vivekanandarajah S, Jeyaseelan E. Pharmacological activities and phytochemical constituents of *Phyllanthus reticulatus* Poir. – A review. *Vingnanam Journal of Science, University of Jaffna*. 2020;15:7–14.
- Vivekanandarajah S, Rajamanoharan PRS. Medicinal values of a food plant – *Limonia acidissima* Groff. *Black Sea Journal of Health Science*. 2021;4:1–6. doi:10.19127/bshealthscience.861779.
- Vivekanandarajah S, Rajamanoharan PRS, Heinrich M. Siddha Medicine in Eastern Sri Lanka Today–Continuity and Change in the Treatment of Diabetes. *Frontiers in Pharmacology*. 2018;9:1022. doi:10.3389/fphar.2018.01022.
- Vivekanandarajah S, Rajamanoharan PRS, Munday M. Plants used to treat diabetes in Sri Lankan Siddha Medicine – An ethnopharmacological review of historical and modern sources. *Journal of Ethnopharmacology*. 2016;198:531–599. doi:10.1016/j.jep.2016.03.060.
- Vivekanandarajah S, Rajamanoharan PRS, Munday M, Heinrich M. Plants currently used to treat diabetes in Sri Lankan Siddha Medicine – an ethnobotanical survey in the Eastern Province.

Presented at: The World Congress Integrative Medicine & Health 2017: Part Three; 2017. Springer Nature, Berlin, Germany. doi:10.1186/s12906-017-1784-2.

Vivekanandarajah S, Rajamanoharan PRS, Heinrich M, Munday M. Preparations and plants used to treat diabetes in Sri Lankan Siddha Medicine. Presented at: The 3rd International Conference on Ayurveda, Unani, Siddha and Traditional Medicine; 2015. Institute of Indigenous Medicine, University of Colombo, Colombo; 67.

Vivekanandarajah S. Antidiabetic plants used by Sri Lankan Tamils. Jaffna Science Association Newsletter. 2021;28(3):3.

Vivekanandarajah S. Plants historically used by Sri Lankan Tamils to treat diabetes. Presented at: The International Dr. Safiye Ali Multidisciplinary Studies Congress in Health Sciences; 2021. İzmir, Turkey.

Watanabe M, Gangitano E, Francomano D, Addressi E, Toscano R, Costantini D, Tuccinardi D, Mariani S, Basciani S, Spera G, Gnassi L, Lubrano C. Mangosteen extract shows a potent insulin sensitizing effect in obese female patients: A prospective randomized controlled pilot study. *Nutrients*. 2018;10:586. doi:10.3390/nu10050586.

Wathsara HPT, Weeratunge HD, Mubarak MNA, Godakumbura PI, Ranasinghe P. In vitro antioxidant and antidiabetic potentials of *Syzygium caryophyllatum* L. Alston. *Evidence-Based Complementary and Alternative Medicine*. 2020. doi:10.1155/2020/9529042.

WebMD LLC. Available at: <https://www.webmd.com/diabetes/default.htm> [Accessed 02 June 2023].

Wedick NM, Pan A, Cassidy A, Rimm EB, Sampson L, Rosner B, Willett W, Hu FB, Sun Q, van Dam RM. Dietary flavonoid intakes and risk of type 2 diabetes in US men and women. *American Journal of Clinical Nutrition*. 2012;95:925–933.

World Health Organization. Monographs on Selected Medicinal Plants, Vol. 1. Available at: <http://apps.who.int/medicinedocs/en/d/Js2200e/> [Accessed 04 August 2023].

World Health Organization. Monographs on Selected Medicinal Plants, Vol. 2. Available at: <http://apps.who.int/medicinedocs/en/d/Js4927e/> [Accessed 04 August 2023].

World Health Organization. Monographs on Selected Medicinal Plants, Vol. 3. Available at: <http://apps.who.int/medicinedocs/en/m/abstract/Js14213e/> [Accessed 04 August 2023].

World Health Organization. Monographs on Selected Medicinal Plants, Vol. 4. Available at: <http://apps.who.int/medicinedocs/en/m/abstract/Js16713e/> [Accessed 04 June 2023].

World Health Organization. Diabetes. Available at: <http://www.who.int/news-room/fact-sheets/detail/diabetes> [Accessed 20 May 2023].

World Health Organization. Available at: https://www.who.int/health-topics/diabetes#tab=tab_1 [Accessed 21 May 2023].

Xi B, Li S, Liu Z, Tian H, Yin X, Huai P, Tang W, Zhou D, Steffen LM. Intake of fruit juice and incidence of type 2 diabetes: A systematic review and meta-analysis. *PLOS One*. 2014;9(3):e93471. doi:10.1371/journal.pone.0093471.

Zulaikha S, Ghafar A, Mediani A, Ramli NS, Abas F. Antioxidant, α -glucosidase, and nitric oxide inhibitory activities of *Phyllanthus acidus* and LC-MS/MS profile of the active extract. *Food Bioscience*. 2018. doi:10.1016/j.fbio.2018.08.009.

Copyright © 2024 The Author(s). This is an open-access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.